AD	

GRANT NO: DAMD17-94-J-4307

TITLE: Genetic Alterations in Familial Breast Cancer: Mapping and Cloning Genes Other than BRCAl

PRINCIPAL INVESTIGATOR: Mary-Claire King, Ph.D.

CONTRACTING ORGANIZATION: University of California
Berkeley, California 94720

REPORT DATE: 13 Sep 95

TYPE OF REPORT: Annual



PREPARED FOR: U.S. Army Medical Research and Materiel

Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;

distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden. to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blan		3. REPORT TYPE AND								
	13 Sep 95	Annual 15 Aug 9								
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS							
Genetic Alterations in Cloning Genes Other th	DAMD17-94-J-4307									
6. AUTHOR(S)										
Mary-Claire King, Ph.D.										
7. PERFORMING ORGANIZATION N	8. PERFORMING ORGANIZATION									
University of Californ		REPORT NUMBER								
Berkeley, California 94720										
a SPONSORING/MONITORING AG	FNCY NAMF(S) AND ADDRESS(ES	1	10. SPONSORING / MONITORING							
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012										
11. SUPPLEMENTARY NOTES		A								
,		•								
12a. DISTRIBUTION / AVAILABILITY	STATEMENT		12b. DISTRIBUTION CODE							
Approved for public release; distribution unlimited										
13. ABSTRACT (Maximum 200 word	de)	A LUCIO								
The purpose of this project is to identify genes other than BRCAl responsible for inherited predisposition to breast cancer, to identify critical mutations, to evaluate the frequency of inherited mutations in these genes among breast cancer patients from the general population, and to determine the role of somatic mutation in these genes in malignant breast tumors. The patient materials are kindreds with at least four cases of breast cancer, in whom disease is not linked to BRCAl. Twenty— one families have been sampled, lymphocyte lines established, and linkage evaluated at multiple chromosomal locales. Linkage analysis of these kindreds reveals convincing linkage to BRCA2 in three families and to the estrogen receptor in three other families. Five other candidate chromosomal regions have been screened as well. Probands from 139 other high—risk kindreds have been contacted. Of these families, 21 will be informative for linkage. Sampling of these 21 extended families is in progress. This will bring the cohort to 42 families in all.										
14. SUBJECT TERMS			15. NUMBER OF PAGES							
BREAST CANCER; GEN	NETICS		14 16. PRICE CODE							
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICA OF ABSTRACT	ATION 20. LIMITATION OF ABSTRACT							
	Unclassified	Unclassified	Unlimited							

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to stay within the lines to meet optical scanning requirements.

- Block 1. Agency Use Only (Leave blank).
- Block 2. Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.
- Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 30 Jun 88).
- Block 4. <u>Title and Subtitle</u>. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.
- Block 5. <u>Funding Numbers</u>. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract PR - Project G - Grant TA - Task

PE - Program WU - Work Unit
Element Accession No.

Block 6. Author(s). Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

- **Block 7.** <u>Performing Organization Name(s) and Address(es)</u>. Self-explanatory.
- **Block 8.** <u>Performing Organization Report Number</u>. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.
- Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.
- **Block 10.** Sponsoring/Monitoring Agency Report Number. (If known)

Block 11. Supplementary Notes. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. <u>Distribution/Availability Statement</u>. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

Block 12b. <u>Distribution Code</u>.

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank. NTIS - Leave blank.

- **Block 13.** Abstract. Include a brief (Maximum 200 words) factual summary of the most significant information contained in the report.
- **Block 14.** <u>Subject Terms</u>. Keywords or phrases identifying major subjects in the report.
- **Block 15.** <u>Number of Pages</u>. Enter the total number of pages.
- **Block 16.** <u>Price Code</u>. Enter appropriate price code (NTIS only).
- Blocks 17. 19. Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.
- Block 20. <u>Limitation of Abstract</u>. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

 χ Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

X Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

 $\underline{\chi}$ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

 $\frac{\chi}{\text{the investigator(s)}}$ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Access	ion For	
ETIS	QRA&I	I
DTIC 3	CAB	
	besser	
Justin	fication_	
	ibution/	
-	lability	
	Avail an	
Dist	Spec	1
1.1		

Mary Clave Kung	8/14/95
PI - Signature	Date

Genetic alterations in familial breast cancer: Mapping and cloning genes other than BRCA1

Annual report for the period August 15, 1994-August 14, 1995

Table of Contents

Front cover	1
SF 298	. 2
Foreword	3
Table of contents	4
Introduction	5
Body of report	7
Conclusions	10
References	11

Introduction

Breast cancer is the most common malignancy among women, with a cumulative risk of 12.6%, or one in eight, by age 85 for a girl born in 1990 (American Cancer Society 1994; California Department Health Services 1994). The existence of a gene or genes responsible for inherited predisposition to breast and ovarian cancer was suggested more than a century ago (Broca 1886) and supported by a vast epidemiological literature for the past 70 years (e.g. Lane-Clayton 1926; Jacobsen 1946; Penrose et al. 1948; Macklin 1959; Anderson 1972; Bain et al. 1980; Schwartz et al. 1985; Newman et al. 1988; Claus et al. 1991). Segregation analysis of a population-based series of families (not selected for family history) indicated that highly penetrant autosomal dominant susceptibility genes were responsible for 5 to 10% of all breast cancer (Newman et al. 1988; Claus et al. 1991) and ovarian cancer (Schildkraut et al. 1989), and that approximately 1 in 200 women in the general population would develop breast cancer as the consequence of a predisposing mutation in one of these genes (Newman et al. 1988; King et al. 1993).

The existence of one gene predisposing to breast and ovarian cancer, BRCA1, was proven by linkage analysis five years ago (Hall et al. 1990) and confirmed with odds >10²⁶:1 (Narod et al. 1991; Easton et al. 1993). In the high-risk families evaluated for linkage, women who inherited a BRCA1 mutation had >80% lifetime risk of breast cancer and were at increased risk for ovarian cancer (Newman et al. 1988; Easton et al. 1993; Ford et al. 1994). BRCA1 was recently isolated by positional cloning (Miki et al. 1994; Futreal et al. 1994; Castilla et al. 1994; Friedman et al. 1994; Simard et al. 1994).

BRCA2, a second breast cancer susceptibility gene, has been mapped to chromosome 13q12 (Wooster et al. 1994). Families with breast cancer linked to BRCA2 are distinguished by a high incidence of male breast cancer. Risk of breast cancer among males predicted from linkage analysis to carry BRCA2 mutations is 6% by age 70; inherited mutations in BRCA2 may be involved in 15% of all male breast cancer. Risks of female breast cancer are similar for BRCA1 and BRCA2. Both ovarian and prostatic cancer, and possibly ocular melanoma, are also at increased frequency in BRCA2 families (D Goldgar, personal communication).

BRCA1 is responsible for a higher proportion of inherited breast and ovarian cancer than is BRCA2, at least in the populations studied so far.

Among approximately 200 families with at least 4 cases of breast cancer, evaluated in various laboratories, ~50% of families have mutations in, and/or convincing linkage to, BRCA1; ~30% appear linked to BRCA2; and ~20% are not (yet) explained by either BRCA1 or BRCA2. In the subset of these families with both ovarian and breast cancer, 75% are attributable to BRCA1, 23% to BRCA2, and only one family appears as-yet-unexplained. Of course, some "unexplained" families may actually carry BRCA1 and/or BRCA2 mutations, but appear unlinked to either locus because the number of noninherited cases (i.e. phenocopies) is high. Other "unexplained" families may reflect another BRCA locus. In our series, breast cancer in two extended families not linked to BRCA1 or BRCA2 is coinherited with the estrogen receptor (Zuppan et al. 1991; see also Body of Report). The total lod score is 3.7 (1.8 and 1.9 for the two families), but no functionally significant mutations in the estrogen receptor coding sequence have been found thus far in either family.

At least two other genes--P53 and the androgen receptor--are also responsible for inherited predisposition to breast cancer in families. Mutations in P53 lead to multiple cancers in families with Li-Fraumeni syndrome, including breast, childhood leukemia, brain, and sarcoma (Malkin et al. 1990). Mutations in the X-linked androgen receptor lead to breast cancer among men with the rare Reifenstein syndrome (Wooster et al. 1992).

Breast cancer risk may also be influenced by more common alleles of other loci conferring moderate risk. Epidemiologic studies have suggested that carriers of mutations in the ataxia telangiectasia (AT) gene are at increased risk of breast cancer (Swift et al. 1991; Borreson et al. 1990), and the proportion of breast cancer attributable to AT carrier status is estimated at 3.8% (Easton 1994). Now that AT has been cloned (Savitsky et al. 1995), it will be possible to test this hypothesis directly.

Other epidemiologic studies suggest that inherited mutations in the HRAS1 minisatellite locus are associated with increased risk of breast cancer, as well as of other common cancers. Inherited mutations in the HRAS1 minisatellite lead to a large number of individually rare alleles, each derived from one of four common progenitors. Although the relative risk of cancer associated with carrying a rare allele at this locus is only 2.0, the aggregate frequency of rare alleles leads to an attributable risk of ~9% of breast cancer in the population as a whole (Krontiris et al. 1993). The mechanism underlying the increased risk remains unknown.

Body of report

High-risk breast cancer families were drawn from two sources. The first series are 20 families from our ongoing linkage studies. In these families, four or more relatives developed breast cancer, but breast cancer was not linked to BRCA1 and no mutations in BRCA1 were detected in the families. The breast cancers present in each of these families are shown in Table 1. Linkage results for BRCA2, the estrogen receptor, chromosome 3p, 8p12-q12, 11q23, 12q, and 16q24 are shown in Table 2.

Table 1. DAMD17-94-J-4307: Characteristics of families with 4+ breast cancers not linked to BRCA1

Family		Breast cancers							
_	All breast	Female < 50	Female > 50	Male breast	fallopian tube				
8	4	1	3						
9	14	6	8						
10	5	3	2						
11	7	3	4						
13	7	4	3	•					
15	5	. 2	3						
16	8	3		5					
17	4	2	2						
18	5	1	4						
19	9	5	3	1					
20	6	1	5						
22	8	1	7		•				
23	5	2	3						
26	11	7	4		1				
33	5	2	3		1				
65	5	4	1						
68	5	2	3						
87	5	4	1		2				
90	9	8	1						
91	7	2	5						
94	5	3	2						
99	6	2	3	1	4				

Table 2. DAMD17-94-J-4307: Linkage results on candidate chromosomes for breast cancer families not linked to BRCA1

16q24	possible	not applicable	not linked	not linked			not applicable		linked		٠		possible			possible	possible		not linked		not applicable
<u>12a</u>	not linked	not applicable	not linked	linked	not linked	not linked	not applicable		possible	not linked	possible	not linked	not linked				not linked	not linked	not linked		not applicable
11q23	not linked	not applicable	not linked	not linked	not linked	not linked	not applicable		not linked	not linked	not linked	not linked	not linked			possible	not linked	not linked	not linked		not applicable
8p12-q12		not applicable	not linked	possible			not applicable					not linked							not linked		not applicable
30	·	not applicable	not linked	not linked			not applicable					not linked							not linked		not applicable
ESR	possible	not linked	not linked	not linked	linked (1.92)	not linked	not applicable	not linked	not linked	not linked	possible	linked (1.85)	not linked	possible	not linked	not linked	uninformative		not linked		not applicable
BRCA2	linked (<1.0)	linked (1.40)	not linked	not linked	not linked	not linked	linked (1.84)	recheck	not linked	not linked	not linked	not linked	not linked	not linked	not linked	not linked	not linked	not linked	not linked	linked (<1.0)	linked (1.09)
Family	80	6	10	_	13	15	16	17	18	19	50	22	23	56	33		89	87	90	94	66

A second series of families has also been developed, based on probands who reported four or more living women in their families with breast cancer. Each participant signed informed consent prior to interview or sampling by our lab.

Each participant provided names, relationships, cancer sites, and ages at cancer diagnosis of their affected relatives. Probands were asked to provide samples of their blood, and affected family members were contacted and invited to participate. In order to contact family members, probands released their relatives' contact information. Often the probands first contacted their relatives, then after permission was given, they released the contact information to our interviewers. Our lab then contacted the relatives, by letter and then by telephone. The relatives were informed of the study by the researchers in a similar manner to the probands. If the relatives were able and willing to participate, they were sent blood collection kits identical to the kits sent to the probands.

Each woman was asked to provide 35 ml blood. Sample collection involved either sampling in person by our staff or sending blood collection kits to the participant being sampled. Each blood collection kit contained a letter from the researchers outlining the goals of the study, a list of rights of medical research subjects, procedures for drawing and packing blood samples, a sheet to document the time and place of the blood draw, a consent form for the study, and a permission form for the release of pathology reports. Upon receipt of the blood samples, lymphocyte lines were established for each individual.

The following are the results of contacting the 139 women from families with four or more affected relatives who completed the original questionnaire:

- 21 blood sample collected from proband; linkage family
- 28 blood sample collected; family not informative or unavailable
- 67 family not informative or not available; no blood sampled
- 16 proband refused or unable to participate
- 5 unable to locate
- <u>2</u> deceased

139 total

When these 21 high-risk families are sampled, the total cohort of high-risk families will number 42.

Conclusions

The following activities will be carried out in the following year:

1. Families linked to BRCA2 will be evaluated for informative meiotic recombination events. When BRCA2 is cloned, mutations in these families will be screened.

2. The estrogen receptor gene will be screened for mutations in the families linked to ESR.

3. Sampling will be completed on the additional linkage families and cell lines established. Families will be screened for linkage to BRCA1. Families in that group will be included in studies of BRCA1 mutations (supported elsewhere) and excluded from this series.

4. Linkage analyses will continue for the candidate chromosomal regions reported in table 2, both for the families listed in tables 1 and 2

and for the new series.

References

American Cancer Society. <u>Cancer Facts and Figures</u>. (1994) American Cancer Society, Atlanta.

Anderson DE. A genetic study of human breast cancer. (1972) J Natl Cancer Inst 28:1500-1504.

Bain C, Speizer FE, Rosner B, Belanger C, Hennekens CH. Family history of breast cancer as a risk indicator for the disease. (1980) Amer J Epidemiol 111:301-308.

Borresen AL; Andersen TI; Tretli S; Heiberg A; Moller P. Breast cancer and other cancers in Norwegian families with ataxia-telangiectasia. (1990) Genes Chrom Cancer 2: 339-40.

Broca PO. Traite des Tumeurs, Vol 1. Paris: P Asselin, 1886-1889.

Brown MA, Nicolai H, Xu C-F, Griffiths BL, Jones KA, Solomon E, Hosking L, Trowsdale J, Black DM, McFarlane R. Regulation of BRCA1. (1994) Nature 372: 733

California Department Health Services. Cancer incidence and mortality in California by detailed race/ethnicity, 1988-1992. (1994) California Department Health Services, Sacramento.

Castilla LH, Couch FJ, Erdos MR, Hoskins KF, Calzone K, Garber JE, Boyd J, Lubin MB, Deshano ML, Brody LC, Collins FS, Weber BL. Mutations in the BRCA1 gene in families with early-onset breast and ovarian cancer. (1994) Nature Genet 8: 387-391

Claus EB, Risch N, Thompson WD. Genetic analysis of breast cancer in the cancer and steroid hormone study. (1991) Amer J Hum Genet 48: 232-242

Easton D, Ford D, Peto J. Inherited susceptibility to breast cancer. (1993) Cancer Surv 18: 95-113

Easton DF, Bishop DT, Ford D, Crockford GP and Breast Cancer Linkage Consortium. Genetic linkage analysis in familial breast and ovarian cancer: results from 214 families. (1993) Amer J Hum Genet 52: 678-701

Easton DF. Cancer risks in A-T heterozygotes. (1994) International Journal of Radiation Biology 66: S177-82.

Ford D, Easton DF, Bishop DT, Narod SA, Goldgar DE and Breast Cancer Linkage Consortium. Risks of cancer in BRCA1-mutation carriers. (1994) Lancet 343: 692-695

Friedman LS, Ostermeyer EA, Szabo CI, Dowd P, Lynch ED, Rowell SE, King M-C. Confirmation of BRCA1 by analysis of germline mutations linked to breast and ovarian cancer in ten families. (1994) Nature Genet 8: 399-404

Futreal PA, Liu Q, Shattuck-Eidens D, Cochran C, Harshman K, Tavtigian S, Bennett LM, Haugen-Strano A, Swensen J, Miki Y, Eddington K, McClure M, Frye C, Weaver-Feldhaus J, Ding W, Gholami Z, Soderkvist P, Terry L, et al. BRCA1 Mutations in primary breast and ovarian cacinomas. (1994) Science 266: 120-122

Hall JM, Lee MK, Newman B, Morrow JE, Anderson LA, Huey B, King M-C (1990) Linkage of early-onset familial breast cancer to chromosome 17q21. (1990) Science 250: 1684-1689

Jacobsen O. Heredity in breast cancer, a genetic and clinical study of 200 probands. (1946) Op dom Biol hered hum, kbh 11: 1-306.

King M-C, Rowell SE, Love SM. Inherited breast and ovarian cancer: What are the risks? What are the choices? (1993) J Amer Med Assoc 269: 1975-1980.

Krontiris TG; Devlin B; Karp DD; Robert NJ; Risch N. An association between the risk of cancer and mutations in the HRAS1 minisatellite locus. (1993) New England Journal of Medicine 329: 517-23.

Lane-Clayton JE. A further report on cancer of the breast with special reference to its associated antecedent conditions. (1926) Reports on Public Health and Medical Subjects, no. 32. Ministry of Health, London.

Macklin MT. Comparison of the number of breast cancer deaths observed in relatives of breast cancer patients and the number expected on the basis of mortality rates. (1959) J Natl Cancer Inst 22: 927-951.

Malkin D, Li FP, Strong LC, Fraumeni JF Jr, Nelson CE, Kim DH, Kassel J, Gryka MA, Bischoff FZ, Tainsky MA, Friend SH. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. (1990) Science 250: 1233-1238.

Miki Y, Swensen J, Shattuck-Eidens D, Futreal PA, Harshman K, Tavtigian S, Liu Q, Cochran C, Bennett LM, Ding W, Bell R, Rosenthal J, Hussey C, Tran T, McClure M, Frye C, Hattier T, Phelps R, Haugen-Strano A, Katcher H, Yakumo K, Gholami Z, Shaffer D, Stone S, Bayer S, Wray C, Bogden R, Dayananth P, Ward J, Tonin P, Narod S, Bristow PK, Norris FH, Helvering L, Morrison P, Rosteck P, Lai M, Barrett JC, Lewis C, Neuhausen S, Cannon-Albright L, Goldgar D, Wiseman R, Kamb A, Skolnick MH. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. (1994) Science; 266:66-71.

Narod SA, Feunteun J, Lynch HT, Watson P, Conway T, Lynch J, Lenoir GM. Familial breast-ovarian cancer locus on chromosome 17q12-q23. (1991) Lancet 338: 82-83

Newman B, Austin MA, Lee M, King M-C (1988) Inheritance of human breast cancer: Evidence for autosomal dominant transmission in high-risk families. (1988) Proc Nat Acad Sci USA 85: 3044-3048

Penrose LS, Mackenzie HJ, Karn MN. A genetical study of human mammary cancer. (1948) Ann Eugen 14: 234-266.

Savitsky K, Bar-Shira A, Gilad S, Rotman G, Ziv Y, Vangaite L, Tagle DA, Smith S, Uziel T, Sfez S, Ashkenazi M, Pecker I, Frydman M, Hamik R, Patanjali SR, Simmons A, Clines G, Sartiel A, Gatti R, Chessa L, Sanal O, Lavin MF, Jaspers NGF, Taylor A, Malcolm R, Arlett CF, Miki T, Weissman SM, Lovett M, Collins FS, Shiloh Y. A single ataxia telangiectasia gene with a product similar to PI-3 kinase. (1995) Science 268:1749-

Schildkraut JM, Risch N, Thompson WD. Evaluating genetic association among ovarian, breast, and endometrial cancer: Evidence for a breast/ovarian cancer relationship. (1989) Amer J Hum Genet 45:521-529.

Schwartz AG, King MC, Belle SH, Satariano WA, Swanson GM. Risk of breast cancer to relatives of young breast cancer patients. (1985) J Natl Cancer Inst 75:665-668.

Simard J, Tonin P, Durocher F, Morgan K, Rommens J, Gingras S, Samson C, Leblanc J-F, Belanger C, Dion F, Liu Q, Skolnick M, Goldar D, Shattuck-Eidens D, Labrie F, Narod SA. Common origins of BRCA1 mutations in Canadian breast and ovarian cancer families. (1994) Nature Genet 8: 392-398

Swift M; Morrell D; Massey RB; Chase CL. Incidence of cancer in 161 families affected by ataxia-telangiectasia. (1991) New Engl J Med 325: 1831-1836.

Wooster R, Mangion J, Eeles R, Smith S, Dowsett M, Averill D, Barrett-Lee P, Easton DF, Ponder BA, Stratton MR. A germline mutation in the androgen receptor gene in two brothers with breast cancer and Reifenstein syndrome. (1992) Nature Genet 2:132-134.

Wooster R., Neuhausen SL, Mangion J, Quirk Y, Ford D, Collins N, Nguyen K, Seal S, Tran T, Averill D, Fields P, Marshall G, Narod S, Lenoir G, Lynch H, Feunteun J, Devilee P, Cornelisse CJ, Menko FH, Daly PA, Ormiston W, McManus R, Pye C, Lewis CM, Cannon-Albright LA, Peto J, Ponder BAJ, Skolnick MH, Easton DF, Goldgar DE, Stratton MR. Localization of a breast cancer susceptibility gene, BRCA2, to chromosome 13q12-13. (1994) Science 265:2088-2090.

Zuppan PJ, Hall JM, Lee MK, Ponglikitmongkol M, King M-C. Possible linkage of the estrogen receptor gene to breast cancer in a family with late-onset disease. (1991) Amer J Human Genet 48:1065-1068.